For Research Use Only

CoraLite® Plus 488-conjugated PSAP Polyclonal antibody



Catalog Number: CL488-10801

Featured Product

Basic Information

Catalog Number:

GenBank Accession Number:

Antigen affinity purification

CL488-10801 Size:

GeneID (NCBI):

Recommended Dilutions:

1000 ug/ml

5660

BC001503

IF/ICC 1:50-1:500

Purification Method:

493 nm / 522 nm

Source:

UNIPROT ID: P07602

Excitation/Emission maxima wavelengths:

Rabbit Isotype:

AG1194

Full Name: prosaposin

Immunogen Catalog Number:

58 kDa

Calculated MW:

Observed MW:

60 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

IF/ICC: HeLa cells, A431 cells

Positive Controls:

Species Specificity:

human

Background Information

The PSAP gene encodes prosaposin, a precursor of four small nonenzymatic glycoproteins termed 'sphingolipid activator proteins' (SAPs) that assist in the lysosomal hydrolysis of sphingolipids. After proteolytic processing of the presaposin protein, these 4 released polypeptides are functional activators. Saposin A is encoded by residues 60 to 143 of PSAP, saposin B by 195 to 275, saposin C by 311 to 390, and saposin D by 405 to 487. There are four 12-14 kDa heatstable glycoproteins. Saposins A-D localize primarily to the lysosomal compartment where they facilitate the catabolism of glycosphingolipids with short oligosaccharide groups. Saposins A-D are required for the hydrolysis of certain sphingolipids by specific lysosomal hydrolases. (PMID: 2001789) Defects in PSAP are the cause of Gaucher disease, Tay-Sachs disease, and metachromatic leukodystrophy (PubMed: 2060627, PMID: 15773042). This PSAP antibody (10801-1-AP) is expected to recognize both saposin A and B.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite® Plus 488 PSAP antibody (CL488-10801) at dilution of 1:200, CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using Coralite® Plus 488 PSAP antibody (CL488-10801) at dilution of 1:200, CL594-Phalloidin (red).



1x10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488-conjugated PSAP Polyclonal antibody (CL488-10801)(red), or 0.4 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).