For Research Use Only

CoraLite® Plus 488-conjugated VDAC1/2 Polyclonal antibody

Catalog Number: CL488-10866 Featured Product



Basic Information

 Catalog Number:
 GenBank Accession Number:

 CL488-10866
 BC008482

 Source:
 GeneID (NCBI):

 Rabbit
 7416

 Isotype:
 UNIPROT ID:

 IgG
 P21796

Immunogen Catalog Number: Full Name:
AG1144 voltage-dependent anion channel 1

Calculated MW: 31 kDa Observed MW: 31 kDa Purification Method: Antigen affinity purification Recommended Dilutions: IF-P: 1:50-1:500

FC (Intra): 0.80 ug per 10^6 cells in a 100 µl suspension

Excitation/Emission maxima wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF-P, FC (Intra) Species Specificity: human, mouse, rat Positive Controls:

IF-P: mouse liver tissue,
FC (Intra): HepG2 cells,

Background Information

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1. C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This polyclonal antibody raised against full-length human VDAC1 protein can cross react with VDAC2.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using CoraLite® Plus 488 VDAC 1/2 antibody (CL488-10866) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488-conjugated VDAC1/2 Polyclonal antibody (CL488-10866)(red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.