For Research Use Only

CoraLite® Plus 488-conjugated CSN2 Polyclonal antibody



Purification Method:

IF/ICC 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-10969

Featured Product

Basic Information

Catalog Number: CL488-10969

1000 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1415

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse

GenBank Accession Number:

BC012629 GeneID (NCBI): 9318 **UNIPROT ID:** P61201 Full Name:

COP9 constitutive photomorphogenic homolog subunit 2 (Arabidopsis)

Calculated MW: 52 kDa Observed MW:

52 kDa

Positive Controls:

IF/ICC : HeLa cells,

Background Information

COPS2 is an essential component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. (refer to UniProt) Catalog#10969-2-AP is a rabbit polyclonal antibody raised against the full-length of human COPS2. The MW of this protein is 52 kDa, and this antibody specially recognises the 52 kDa

Storage

Applications


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 CSN2 antibody (CL488-10969) at dilution of 1:200, CL594-Phalloidin (red).



1x10^6 HeLa cells were intracellularly stained with 0.8 ug Coralite® Plus 488 Csn2 Polyclonal Antibody (CL488-10969)(red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit 1gG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).