For Research Use Only

## CoraLite® Plus 488-conjugated IRF3 Polyclonal antibody

Catalog Number: CL488-11312

Featured Product

1 Publications



**Basic Information** 

Catalog Number: CL488-11312

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1858

GenBank Accession Number:

BC009395
GeneID (NCBI):
3661
UNIPROT ID:
Q14653
Full Name:

interferon regulatory factor 3

Calculated MW: 47 kDa Observed MW: 50-60 kDa Purification Method:

Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:50-1:500 FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

Tested Applications: IF/ICC, FC (Intra) Cited Applications:

IF

Species Specificity:

human
Cited Species:
human

Positive Controls:

IF/ICC: HepG2 cells,

FC (Intra): human peripheral blood lymphocytes,

## **Background Information**

The virul-induced expression of interferon(IFN) genes in infected cells implicate in the interplay of several constitutively expressed and virus-activated transcription factors. A family of IFN regulatory factors(IRFs) have been shown to has a role in the transcription of IFN genes as well as IFN-stimulated genes. IRF3 is a novel key transcriptional regulator of type I IFN-dependent immune responses and involves in the innate immune response against DNA and RNA viruses, by binding to the promoters of IFN. It located in the cytoplasm of uninfected cells in an inactive form, and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, could be phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization, nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes. IRF3 exists some isoforms with MV 47-49 kDa, 33 kDa and 12-16 kDa.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Lin Zhu	38285487	Elife	IF

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 IRF3 antibody (CL488-11312) at dilution of 1:200.



1x10^6 human peripheral blood lymphocytes were intracellularly stained with 0.8 ug CoraLite® Plus 488-conjugated IRF3 Polyclonal antibody (CL488-11312)(red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-3000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.