For Research Use Only

CoraLite® Plus 488-conjugated ASGR1 Polyclonal antibody



Catalog Number: CL488-11739

Featured Product

1 Publications

GeneID (NCBI):

UNIPROT ID:

Full Name:

Calculated MW:

P07306

GenBank Accession Number:

asialoglycoprotein receptor 1

Basic Information

Catalog Number:

CL488-11739 BC032130

1000 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG2310

291 aa, 33 kDa

IF-P: rat liver tissue, human liver cancer tissue

Purification Method:

IF-P 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Applications

Tested Applications:

IF-P, FC

Cited Applications:

IF

Species Specificity: human, mouse, rat Cited Species:

mouse

Background Information

Asialoglycoprotein receptor (ASGPR), also known as the hepatic galactose/N-acetylglucosamine (GlcNAc) receptor or Ashwell receptor, is a C-type lectin expressed exclusively in hepatic parenchymal cells. ASGPR consists of two subunits, a major subunit (ASGR1, HL-1) and a minor subunit (ASGR2, HL-2), and specifically recognizes terminal βlinked galactose or GlcNAc on circulating glycoproteins or cells. This receptor plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins that contain terminal galactose or GlcNAc residues. ASGPR may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery. This antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

Notable Publications


Author Application **Pubmed ID** Journal Tian Tian MedComm (2020) 37829506

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite® Plus 488 ASGR1 antibody (CL488-11739) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed rat liver tissue using CoraLite® Plus 488 ASGR1 antibody (CL488-11739) at dilution of 1:200.



1X10^6 HepG2 cells were surface stained with 0.4 ug CoraLite® Plus 488 Anti-Human ASGR1 (CL488-11739) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit 1gG control Rabbit PolyAb (CL488-30000, Clone:) (blue). Cells were not fixed.