## For Research Use Only

## CoraLite® Plus 488-conjugated CREB1 Polyclonal antibody

Catalog Number:CL488-12208

Featured Product


## Antibodies | ELISA kits | Proteins WWW.ptglab.com

Basic Information	Catalog Number: CL488-12208	GenBank Accession Number: BC010636	Purification Method: Antigen affinity purification	
	Size: 1000 µg/ml	GenelD (NCBI): 1385	Recommended Dilutions: IF/ICC 1:50-1:500	
	Source: Rabbit	UNIPROT ID: P16220 Full Name: cAMP responsive element binding protein 1	Excitation/Emission maxima wavelengths: 493 nm / 522 nm	
	Isotype: IgG Immunogen Catalog Number:			
	AG2852	Calculated MW: 341 aa, 35 kDa		
		Observed MW: 43-46 kDa		
Applications	Tested Applications:	Tested Applications:Positive Controls:IF/ICC, FC (Intra)IF/ICC : HeLa cells,Species Specificity:IF/ICC : HeLa cells,human, mouse, rat, monkeyIF/ICC : HeLa cells,		
	Species Specificity:			
Background Informatio	CREB1, also named as CREB, belongs to the bZIP family, containing one bZIP domain and one KID (kinase-inducible) domain. This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. This protein is stimulated by phosphorylation. Phosphorylation of both Ser-133 and Ser-142 in the SCN regulates the activity of CREB and participates in circadian rhythm generation. Phosphorylation of Ser-133 allows CREBBP binding. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. CREB1 is sumoylated by SUMO1. Sumoylation on Lys-304, but not on Lys-285, is required for nuclear localization of this protein. Sumoylation is enhanced under hypoxia, promoting nuclear localization and stabilization. Defects in CREB1 may be a cause of angiomatoid fibrous histiocytoma (AFH), a distinct variant of malignant fibrous histiocytoma that typically occurs in children and adolescents and is manifest by nodular subcutaneous growth. A chromosomal aberration involving CREB1 is found in a patient with angiomatoid fibrous histiocytoma. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type. CREB1 exists some isoforms and range of calculated molecular weight of isoforms are 35-37 kDa and 25 kDa, but the modified CREB1 protein is about 43 kDa (PMID: 25883219 ).			
	rhythm generation. Phosphorylati TORC coactivators which act inder Sumoylation on Lys-304, but not c enhanced under hypoxia, promoti angiomatoid fibrous histiocytoma children and adolescents and is m CREB1 is found in a patient with a generates a EWSR1/CREB1 fusion some isoforms and range of calcu	on of Ser-133 allows CREBBP binding. To pendently of Ser-133 phosphorylation. O on Lys-285, is required for nuclear localiz ng nuclear localization and stabilization a (AFH), a distinct variant of malignant f aanifest by nodular subcutaneous growth ngiomatoid fibrous histiocytoma. Trans gene that is most common genetic abno lated molecular weight of isoforms are	ranscription activation is enhanced by the REB1 is sumoylated by SUMO 1. ration of this protein. Sumoylation is n. Defects in CREB1 may be a cause of ibrous histiocytoma that typically occurs n. A chromosomal aberration involving location t(2;22)(q33;q12) with CREB1 prmality in this tumor type. CREB1 exists	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite® Plus 488 CREB1 antibody (CL488-12208) at dilution of 1:100.



1X10<sup>^</sup>6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human CREB1 (CL488-12208) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).