For Research Use Only

## CoraLite® Plus 488-conjugated TDP-43 (C-terminal) Polyclonal antibody



Catalog Number: CL488-12892

Featured Product

**Basic Information** 

Catalog Number: CL488-12892

1000 µ g/ml Source: Rabbit

Isotype:

BC001487

GeneID (NCBI):
23435

UNIPROT ID: Q13148 Full Name:

TAR DNA binding protein

GenBank Accession Number:

Calculated MW: 43 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

Tested Applications: FC (Intra), IF/ICC Species Specificity: human, mouse, rat Positive Controls:

IF: HeLa cells,

## **Background Information**

Transactivation response (TAR), DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43), was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major component of the tau-negative and ubiquitin-positive inclusions that characterize amyotrophic lateral sclerosis (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). 12892-1-AP is a rabbit polyclonal antibody raised against the C-terminal amino acids of human TDP-43. This antibody recognizes the cleavage product of 20-30 kDa in addition to the native and phosphorylated forms of TDP-43. Immunohistochemical analyses of TDP-43 using this antibody detect both normal diffuse nuclear staining and insoluble inclusions in pathologic tissues. Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phosphoprotein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (17023659,19823856,21666678,22193176)

Recently TDP-43 has been reported to be overexpressed in triple negative breast cancer (TNBC) and it may be a potential target for TNBC diagnosis and drug design. (29581274)

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Ruffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite® Plus 488 TDP-43 (C-terminal) antibody (CL488-12892) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human TDP-43 (Cterminal) (CL488-12892) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).