For Research Use Only

CoraLite® Plus 488-conjugated TFAP2A,AP-2 Polyclonal antibody



Catalog Number: CL488-13019

Featured Product

Basic Information

Catalog Number: CL488-13019

1000 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG4112

BC017754

GenBank Accession Number: GeneID (NCBI): 7020 **UNIPROT ID:** P05549 Full Name: transcription factor AP-2 alpha

(activating enhancer binding protein

2 alpha) Calculated MW:

431 aa, 47 kDa Observed MW: 47 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF/ICC

Species Specificity:

human

Positive Controls:

IF: HepG2 cells,

Background Information

The activator protein-2 (AP-2) family of transcription factors comprises five 52-kDa isoforms (AP-2 a , AP-2 β , AP- 2γ , AP-2 δ , and AP-2 ϵ), which share a common structure: a proline/glutamine-rich transactivation domain in the N-terminal region and a helix-span-helix domain in the C-terminal region, which mediates dimerization and sitespecific DNA binding. Depending on the cellular context, the AP-2 transcription factors are individually associated either with cell differentiation and development or with cancer progression/regression (PMID: 21966377). TFAP2A (AP-2-alpha) is the only AP-2 protein required for early morphogenesis of the lens vesicle. Together with the CITED2 coactivator, stimulates the PITX2 P1 promoter transcription activation (PMID: 11694877).


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 TFAP2A,AP-2 antibody (CL488-13019) at dilution of 1:200.