For Research Use Only

CoraLite® Plus 488-conjugated PARP1 Polyclonal antibody



Catalog Number: CL488-13371

Featured Product

Basic Information

Catalog Number: CL488-13371

Size: Genel D (NCBI): 1000 ug/ml 142

Source: UNIPROT ID:
Rabbit P09874
Isotype: Full Name:

IgG poly (ADP-ribose) polymerase 1

Immunogen Catalog Number: Calculated MW: 1014 aa, 113 kDa
Observed MW:

Observed MW: 113-116 kDa, 89 kDa

GenBank Accession Number:

BC037545

. .

Purification Method: Antigen affinity purification Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat

Positive Controls:

IF/ICC: HepG2 cells,

Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24-kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered as an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and gave rise to fragments ranging from 42-89-kDa. This antibody was generated against the C-terminal region of human PARP1 and it recognizes the full-length as well as the cleavage of the PARP1.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 PARP1 antibody (CL488-13371) at dilution of 1:200.



1X10^6 K-562 cells were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human PARP1 (CL488-13371) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 PARP1 antibody (CL488-13371) at dilution of 1:200, CL594-Phalloidin (red).