## For Research Use Only

## CoraLite® Plus 488-conjugated XPA Polyclonal antibody

Catalog Number: CL488-16462

| Basic Information      | Catalog Number:<br>CL488-16462  | GenBank Accession Number:<br>BC014965                           | Purification Method:<br>Antigen affinity purification         |  |
|------------------------|---|---|---|--|
|                        | Size:<br>1000 µ g/ml<br>Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG9536   | GeneID (NCBI):<br>7507  | Excitation/Emission maxima<br>wavelengths:<br>493 nm / 522 nm |  |
|                        |   | UNIPROT ID:<br>P23025   |   |  |
|                        |   | Full Name:<br>xeroderma pigmentosum,<br>complementation group A |   |  |
|                        |   | Calculated MW:<br>374aa,42 kDa; 273aa,31 kDa<br>Observed MW:    | :Da   |  |
|                        |   | 40 kDa  |   |  |
| Applications           | Tested Applications:<br>FC (Intra)  |   |   |  |
|                        | Species Specificity:<br>human, mouse  |   |   |  |
| Background Information | XPA, also named as DNA repair protein complementing XP-A cells, is a 272 amino acid protein, which belongs to the XPA family. XPA is expressed in various cell lines and in skin fibroblasts. XPA is involved in DNA excision repair. Initiates repair by binding to damaged sites with various affinities, depending on the photoproduct and the transcriptional state of the region. XPA is required for UV-induced CHEK1 phosphorylation and the recruitment of CEP164 to cyclobutane pyrimidine dimmers (CPD), sites of DNA damage after UV irradiation. The calculated molecular weight of XPA is 31 kDa, but modified XPA protein is about 40 kDa (PMID: 17848622). |   |   |  |
| Storage                | Storage:<br>Store at -20°C. Avoid exposure to light. Stable for one year after shipment.<br>Storage Buffer:<br>PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.<br>Aliquoting is unnecessary for -20°C storage  |   |   |  |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Antibodies | ELISA kits | Proteins

www.ptglab.com

## Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human XPA (CL488-16462) (red), or 0.8 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).