## For Research Use Only

## CoraLite® Plus 488-conjugated Cytokeratin 20 Polyclonal antibody

Catalog Number:CL488-17329




Basic Information	Catalog Number: CL488-17329	GenBank Accession Number: BC031559	Purification Method: Antigen affinity purification
	Size: 1000 ug/ml	GenelD (NCBI): 54474	Recommended Dilutions: IF/ICC 1:50-1:500
	Source: Rabbit	UNIPROT ID: P35900	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
	lsotype: lgG Immunogen Catalog Number: AG11308	Full Name: keratin 20	
		Calculated MW: 424 aa, 48 kDa	
		Observed MW: 46 kDa	
Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : HT-29 cells,	
	Species Specificity: human, mouse, rat, rabbit		
Background Information	Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. Keratin 20 is a type I cytokeratin. It is a major cellular protein of mature enterocytes and goblet cells and is specifically found in the gastric and intestinal mucos		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliguoting is unnecessary for -20°C storage		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HT-29 cells using CoraLite® Plus 488 Cytokeratin 20 antibody (CL488-17329) at dilution of 1:200.



1x10^6 HT-29 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488-conjugated Cytokeratin 20 Polyclonal antibody (CL488-17329) (red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).