## For Research Use Only

## CoraLite® Plus 488-conjugated Caspase 3/P17/P19 Polyclonal antibody



Catalog Number:CL488-19677


Featured Product

<b>Basic Information</b>	Catalog Number: CL488-19677	GenBank Accession Number: NM_004346	Purification Method: Antigen affinity purification	
	Size: 1000 ug/ml	GeneID (NCBI): 836	Recommended Dilutions: IF/ICC 1:50-1:500	
	Source: Rabbit	UNIPROT ID: P42574	Excitation/Emission maxima wavelengths:	
	lsotype: IgG	Full Name: 493 nm / 522 nm   caspase 3, apoptosis-related cysteine   peptidase	493 nm / 522 nm eine	
		Calculated MW: 32 kDa Observed MW: 32-35 kDa, 17 kDa, 19 kDa		
Applications	Tested Applications:	Positive C	ontrols:	
	IF/ICC Species Specificity: human, mouse, rat	IF/ICC : HeLa cells, NIH/3T3 cells		
	Corporation of family of and and	Caspases, a family of endoproteases, are critical players in cell regulatory networks controlling inflammation and cell death. Initiator caspases (caspase-2, -8, -9, -10, -11, and -12) cleave and activate downstream effector caspases (caspase-3, -6, and -7), which in turn execute apoptosis by cleaving targeted cellular proteins. Caspase 3 (also named CPP32, SCA-1, and Apopain) proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at the beginning o apoptosis. Caspase 3 plays a key role in the activation of sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase 3 can also form heterocomplex with other proteins and performs the molecular mass of 50-70 kDa. This antibody can recognize p17, p19 and p32 of Caspase 3.		
Background Informati	cell death. Initiator caspases (caspase-3, -6, and -7), which named CPP32, SCA-1, and Ap apoptosis. Caspase 3 plays a between the basic helix-loop also form heterocomplex wit	(caspase-2, -8, -9, -10, -11, and -12) cleave in turn execute apoptosis by cleaving targ opain) proteolytically cleaves poly(ADP-r key role in the activation of sterol regulat -helix leucine zipper domain and the men n other proteins and performs the molecul	and activate downstream effector caspases eted cellular proteins. Caspase 3 (also ibose) polymerase (PARP) at the beginning o ory element binding proteins (SREBPs) nbrane attachment domain. Caspase 3 can	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using CoraLite® Plus 488 Caspase 3/p17/p19 antibody (CL488-19677) at dilution of 1:100.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 Caspase 3/p17/p19 antibody (CL488-19677) at dilution of 1:200.