For Research Use Only

CoraLite® Plus 488-conjugated GLUT1 Polyclonal antibody



Purification Method:

IF-P 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-21829

Featured Product

Basic Information

Catalog Number: CL488-21829

Size: 1000 µg/ml

Source: Rabbit P11166 Full Name: Isotype:

Immunogen Catalog Number:

AG16282

GenBank Accession Number: BC121804

GeneID (NCBI): **UNIPROT ID:**

solute carrier family 2 (facilitated glucose transporter), member 1

Calculated MW: 492 aa. 54 kDa

Observed MW: 45-55 kDa

Positive Controls:

IF-P: mouse brain tissue,

Applications

Tested Applications: IF-P, FC (Intra) Species Specificity:

human, mouse, rat

Background Information

SLC2A1, also known as GLUT1, is an ubiquitously expressed glucose transporter and responsible for the basal level of glucose uptake in most cell types. Human erythrocytes express the highest level of the SLC2A1. Defects in SLC2A1 are the cause of GLUT1 deficiency syndrome type 1 and type 2. High expression of SLC2A1 has been reported to be a reliable immunohistochemical marker for juvenile hemangiomas. GLUT1 protein may appear as two or more distinct forms among 43 kDa to 55 kDa due to the different glycosylation state. And the conversion of highly glycosylated form of GLUT1 to less glycosylated form has been reported to correlate to differentiation (PMID: $8263524, 23302780).\ 21829-1-AP\ antibody\ can\ also\ detect\ the\ 25\ kDa\ degradation\ protein\ in\ SDS-PAGE$ (PMID:18387950).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using CoraLite® Plus 488 GLUT1 antibody (CL488-21829) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using CoraLite® Plus 488 GLUT1 antibody (CL488-21829) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using CoraLite® Plus 488 GLUT1 antibody (CL488-21829) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 Jurkat cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human GLUT1 (CL488-21829)(red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-3000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).