For Research Use Only

CoraLite® Plus 488-conjugated Angiotensinogen/AGT Polyclonal antibody



Purification Method:

IF/ICC 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-23972

Basic Information

Catalog Number: CL488-23972 Concentration: 1000 ug/ml

Source: U
Rabbit P
Isotype: F
IgG a

Immunogen Catalog Number:

AG21121

IF/ICC, FC (Intra)
Species Specificity:

Tested Applications:

human

GenBank Accession Number:

BC011519 GeneID (NCBI):

UNIPROT ID: P01019 Full Name:

angiotensinogen (serpin peptidase inhibitor, clade A, member 8)

Calculated MW: 485 aa, 53 kDa

Observed MW: 52 kDa

Positive Controls:

IF/ICC: HepG2 cells, HeLa cells

Background Information

Angiotensinogen is a precursor of angiotensin II (Ang II), is expressed and synthesized largely in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. It has a key role in mediating vascular constriction and regulating salt and fluid homeostasis. The resulting product, angiotensin I, is then cleaved by angiotensin converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II. Mutations in this gene are associated with susceptibility to essential hypertension, and can cause renal tubular dysgenesis, a severe disorder of renal tubular development. Defects in this gene have also been associated with non-familial structural atrial fibrillation, and inflammatory bowel disease.

Storage

Applications


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



1x10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human Angiotensinogen (CL488-23972)(red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using CoraLite® Plus 488 Angiotensinogen antibody (CL488-23972) at dilution of 1:200.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 Angiotensinogen antibody (CL488-23972) at dilution of 1:200.