For Research Use Only

CoraLite® Plus 488-conjugated KEAP1 Monoclonal antibody

Catalog Number:CL488-60027

Featured Product

1 Publications





| Basic Information | Catalog Number: CL488-60027 | GenBank Accession Number: BC002930 | Purification Method: Thiophilic affinity chromatograph |
|--|---|---|--|
| | Size: | GenelD (NCBI): | CloneNo.: |
| | 1000 ug/ml Source: | 9817 UNIPROT ID: | 4G10H9 Recommended Dilutions: |
| | Mouse | Q14145 | IF/ICC 1:50-1:500 |
| | Isotype: IgA | Full Name: kelch-like ECH-associated protein 1 | Excitation/Emission maxima wavelengths: |
| | Immunogen Catalog Number: AG0779 | Calculated MW: 624 aa, 70 kDa | 493 nm / 522 nm |
| | | Observed MW: 70 kDa | |
| Applications | Tested Applications: IF/ICC | Positive Controls: IF/ICC : HeLa cells, | |
| | Cited Applications: | | |
| | Species Specificity: human, mouse, rat | | |
| | Cited Species: mouse | | |
| | | | |
| Background Information | ubiquitin ligase, which can ubiqu distinct sites in the N-terminal re and/or its physiological conseque protein for the E3 ubiquitin ligase | A0132 and KLHL19, is part of a multiprotei itinate the N-terminal domain of NRF2. Tv gion of NRF2, the ETGE and DLG sites, whi ences. KEAP1 retains NFE2L2/NRF2 in the complex formed by CUL3 and RBX1. It als antibody raised against residues near the | vo molecules of KEAP1 bind to two th affect the KEAP1-NRF2 interaction sytosol. It functions as substrate adap o retains BPTF in the cytosol. This |
| | ubiquitin ligase, which can ubiqu distinct sites in the N-terminal re- and/or its physiological conseque protein for the E3 ubiquitin ligase antibody is a mouse monoclonal | itinate the N-terminal domain of NRF2. Tv gion of NRF2, the ETGE and DLG sites, whic ences. KEAP1 retains NFE2L2/NRF2 in the c e complex formed by CUL3 and RBX1. It als | vo molecules of KEAP1 bind to two th affect the KEAP1-NRF2 interaction sytosol. It functions as substrate adap o retains BPTF in the cytosol. This |
| Background Information Notable Publications | ubiquitin ligase, which can ubiqu distinct sites in the N-terminal re- and/or its physiological conseque protein for the E3 ubiquitin ligase antibody is a mouse monoclonal | itinate the N-terminal domain of NRF2. Tv gion of NRF2, the ETGE and DLG sites, whice ences. KEAP1 retains NFE2L2/NRF2 in the c e complex formed by CUL3 and RBX1. It als antibody raised against residues near the | vo molecules of KEAP1 bind to two ch affect the KEAP1-NRF2 interaction cytosol. It functions as substrate adap o retains BPTF in the cytosol. This C terminus of human KEAP1. Application |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data







Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 KEAP1 antibody (CL488-60027, Clone: 4G10H9) at dilution of 1:200.