For Research Use Only

CoraLite® Plus 488-conjugated TNFalpha Monoclonal antibody

Catalog Number:CL488-60291 Featured Product


4 Publications




Basic Information	Catalog Number: CL488-60291	GenBank Accession Number: BC028148	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	1000 ug/ml	7124	7B8A11	
	Source: Mouse	ENSEMBL Gene ID: ENSG00000232810	Recommended Dilutions: IF-P 1:50-1:500	
	lsotype: IgG2b	UNIPROT ID: P01375	Excitation/Emission maxima wavelengths:	
	Immunogen Catalog Number: AG11413	Full Name: tumor necrosis factor (TNF superfamily, member 2)	493 nm / 522 nm	
		Calculated MW: 233 aa, 26 kDa		
Applications	Tested Applications:	Positive Controls:		
	IF-P Cited Applications: WB, IHC, IF	IF-P : human breast cancer tissue,		
	Species Specificity: human			
	Cited Species: human, mouse			
	TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved b TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, INS resistance, and cancer. The antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.			
Background Information	TNF-alpha converting enzyme (TA circulation. It is produced chiefly such as CD4+ lymphocytes, NK ce functions through its receptors TN of a wide spectrum of biological p and coagulation. This cytokine ha	ACE) to release the soluble 17 kDa mo by activated macrophages, although i lls, neutrophils, mast cells, eosinophil FRSF1A/TNFR1 and TNFRSF1B/TNFBR processes including cell proliferation, s been implicated in a variety of dise	nomer, which forms homotrimers in t can be produced by many other cell types s, and neurons. It can bind to, and thus . This cytokine is involved in the regulation differentiation, apoptosis, lipid metabolism ases, including autoimmune diseases, INS	
	TNF-alpha converting enzyme (TA circulation. It is produced chiefly such as CD4+ lymphocytes, NK ce functions through its receptors TN of a wide spectrum of biological p and coagulation. This cytokine ha resistance, and cancer. The antibo	ACE) to release the soluble 17 kDa mo by activated macrophages, although i lls, neutrophils, mast cells, eosinophil FRSF1A/TNFR1 and TNFRSF1B/TNFBR processes including cell proliferation, s been implicated in a variety of dise dy is conjugated with CL488, Ex/Em 4	nomer, which forms homotrimers in t can be produced by many other cell types s, and neurons. It can bind to, and thus . This cytokine is involved in the regulation differentiation, apoptosis, lipid metabolism ases, including autoimmune diseases, INS .88 nm/515 nm.	
	TNF-alpha converting enzyme (TA circulation. It is produced chiefly such as CD4+ lymphocytes, NK cel functions through its receptors TN of a wide spectrum of biological p and coagulation. This cytokine ha resistance, and cancer. The antibo	ACE) to release the soluble 17 kDa mo by activated macrophages, although i lls, neutrophils, mast cells, eosinophil FRSF 1A/TNFR1 and TNFRSF 1B/TNFBR processes including cell proliferation, s been implicated in a variety of dise idy is conjugated with CL488, Ex/Em 2 Pubmed ID Journal	nomer, which forms homotrimers in t can be produced by many other cell types s, and neurons. It can bind to, and thus . This cytokine is involved in the regulation differentiation, apoptosis, lipid metabolism ases, including autoimmune diseases, INS .88 nm/515 nm. Application	
	TNF-alpha converting enzyme (TA circulation. It is produced chiefly such as CD4+ lymphocytes, NK cel functions through its receptors TN of a wide spectrum of biological g and coagulation. This cytokine ha resistance, and cancer. The antibo Author Qing Xiong	ACE) to release the soluble 17 kDa mo by activated macrophages, although i lls, neutrophils, mast cells, eosinophil FRSF 1A/TNFR1 and TNFRSF 1B/TNFBF processes including cell proliferation, is been implicated in a variety of dise idy is conjugated with CL488, Ex/Em 2 Pubmed ID Journal 31590127 Biomed Pharmace	nomer, which forms homotrimers in t can be produced by many other cell types s, and neurons. It can bind to, and thus . This cytokine is involved in the regulatior differentiation, apoptosis, lipid metabolism ases, including autoimmune diseases, INS .88 nm/515 nm. Application other WB	
Background Information	TNF-alpha converting enzyme (TA circulation. It is produced chiefly such as CD4+ lymphocytes, NK cel functions through its receptors TN of a wide spectrum of biological p and coagulation. This cytokine ha resistance, and cancer. The antibo Author Qing Xiong Jing Ma	ACE) to release the soluble 17 kDa mo by activated macrophages, although i lls, neutrophils, mast cells, eosinophil FRSF 1A/TNFR1 and TNFRSF 1B/TNFBR processes including cell proliferation, s been implicated in a variety of dise idy is conjugated with CL488, Ex/Em 2 Pubmed ID Journal	nomer, which forms homotrimers in t can be produced by many other cell types s, and neurons. It can bind to, and thus . This cytokine is involved in the regulatior differentiation, apoptosis, lipid metabolism ases, including autoimmune diseases, INS .88 nm/515 nm. Application	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CL488-60291 (TNF alpha antibody) at dilution of 1:50.



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CL488-60291 (TNF alpha antibody) at dilution of 1:50.