For Research Use Only

CoraLite® Plus 488 Anti-Human CD25/IL-2RA (BC96)

Catalog Number: CL488-65096



Basic Information

Catalog Number:

CL488-65096

Size:

100tests, 5 µl/test

Source: Mouse

Isotype: IgG1, kappa GenBank Accession Number: BC156627

GeneID (NCBI):

559

UNIPROT ID: P01589 Full Name:

interleukin 2 receptor, alpha

Calculated MW: 272 aa, 31 kDa Purification Method: Affinity purification

CloneNo.:

BC96

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

Proliferation of T lymphocytes is triggered by the interaction of IL-2 with its specific receptor following T lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4+Foxp3+Treg cells and a subset of human CD4+ memory T cells (PMID: 22585674).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.1% sodium azide and 0.5% BSA, pH 7.3.

Selected Validation Data



1X10^6 human peripheral blood mononuclear cells (PBMCs) were surface stained with APC Anti-Human CD4 (APC-65134, Clone: OKT4) and 5.00 ul Coralite® Plus 488-conjugated Anti-Human CD25 (CL488-65096, Clone: BC96). Cells were not fixed.



1X10^6 human peripheral blood mononuclear cells (PBMCs) were surface stained with APC Anti-Human CD4 (APC-65134, Clone: OKT4) and CoraLite® Plus 488-conjugated Mouse IgG1 isotype control antibody. Cells were not fixed.