For Research Use Only

CoraLite® Plus 488 Anti-Human CD25/IL-2RA (BC96)

Catalog Number:CL488-65096




Basic Information	Catalog Number: CL488-65096	GenBank Accession Number: BC156627	Purification Method: Affinity purification					
	Concentration: 100tests, 5 µl/test	GenelD (NCBI): 3559	CloneNo.: BC96					
	Source: Mouse Isotype: IgG1, kappa	UNIPROT ID: P01589 Full Name: interleukin 2 receptor, alpha Calculated MW: 272 aa, 31 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm					
				Applications	Tested Applications: FC Species Specificity: human		Positive Controls: FC : human PBMCs,	
				Background Information	Proliferation of T lymphocytes is triggered by the interaction of IL-2 with its specific receptor following T lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three differen proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells,			
	activated B cells, some thym	activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4+Foxp3+ Treg cells and a subset of human CD4+ memory T cells (PMID: 22585674).						
Storage	Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide and 0.5% BSA, pH7.3							


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 CD3/CD28 activated (3 day) human PBMCs were surface stained with 5 ul CoraLite® Plus 488 Anti-Human CD25/IL-2RA (CL488-65096, Clone: BC96) (red) or CoraLite® Plus 488 Mouse IgG1 Isotype Control (CL488-65124, Clone: MOPC-21) (blue). 1x10^6 unactivated human PBMCs were surface stained with 5 ul CoraLite® Plus 488 Anti-Human CD25/IL-2RA (CL488-65096, Clone: BC96) (black dashed). Cells were not fixed. Cells were treated with Fc Receptor



1x10^6 CD3/CD28 activated (3 day) human PBMCs were surface stained with 5 ul CoraLite® Plus 488 Anti-Human CD25/IL-2RA (CL488-65096, Clone: BC96) (right) or CoraLite® Plus 488 Mouse IgG1 Isotype Control (CL488-65124, Clone: MOPC-21) (left). 1x10^6 unactivated human PBMCs were surface stained with 5 ul CoraLite® Plus 488 Anti-Human CD25/IL-2RA (CL488-65096, Clone: BC96) (center). Cells were not fixed. Cells were treated with Fc Receptor Block