For Research Use Only

CoraLite® Plus 488 Anti-Human CD27 (0323)



Purification Method:

Affinity purification

Recommended Dilutions:

FC: 5 ul per 10^6 cells in 100 μ l

Excitation/Emission maxima

CloneNo.:

suspension

wavelengths:

493 nm / 522 nm

0323

Catalog Number: CL488-65132

Basic Information

Catalog Number:

CL488-65132 Concentration:

100tests, 5 µl/test

Source: Mouse

lgG1

Isotype:

Full Name:

Calculated MW:

939

Applications

Tested Applications:

Species Specificity:

Human

GenBank Accession Number:

BC012160 GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000139193

UNIPROT ID: P26842

CD27 molecule

29 kDa

Positive Controls:

FC: human peripheral blood lymphocytes,

Background Information

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligation of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1X10^6 human peripheral blood lymphocytes were surface stained with APC-Anti-Human CD19 (APC-65145, Clone: SJ25C1) and 5.00 ul CoraLite® Plus 488-conjugated Anti-Human CD27 (CL488-65132, Clone: O323). Cells were not fixed.



1X10^6 human peripheral blood lymphocytes were surface stained with APC-Anti-Human CD19 (APC-65145, Clone: SJ25C1) and CoraLite® Plus 488-conjugated Mouse IgG1 isotype control antibody. Cells were not fixed.