For Research Use Only

## CoraLite® Plus 488 Anti-Human CD8 (UCHT4)

proteintech®

Antibodies | ELISA kits | Proteins

www.ptglab.com

Catalog Number: CL488-65204

**Basic Information** 

Catalog Number:

CL488-65204

Size:

100tests, 5  $\mu$  l/test

Source: Mouse

Isotype: IgG2a GenBank Accession Number:

BC025715 GeneID (NCBI):

925

ENSEMBL Gene ID: ENSG00000153563

UNIPROT ID: P01732

Full Name: CD8a molecule Calculated MW: 235 aa, 26 kDa Purification Method:

Protein A purification

CloneNo.: UCHT4

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

Human

**Background Information** 

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage:

Store at 2-8 °C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA, pH 7.3.

## **Selected Validation Data**



1X10^6 human peripheral blood lymphocytes were surface stained with APC-Anti-Human CD3 (APC-65151, Clone: UCHT1) and 5 ul Coralite® Plus 488-conjugated Anti-Human CD8 (CL488-65204, Clone: UCHT4). Cells were not fixed.



1X10^6 human peripheral blood lymphocytes were surface stained with APC-Anti-Human CD3 (APC-65151, Clone: UCHT1) and Coralite® Plus 488-conjugated Mouse IgG2a isotype control. Cells were not fixed.