

For Research Use Only

CoraLite® Plus 488 Anti-Human CD83 (HB15e) Mouse IgG2a Recombinant Antibody

Catalog Number: CL488-65581



Basic Information

Catalog Number:	BC030830	GenBank Accession Number:	Purification Method:
CloneNo.:	HB15e	Concentration:	Protein A purification
Source:	Mouse	GenID (NCBI):	100 tests, 5 µl/test
Isotype:	IgG2a	ENSEMBL Gene ID:	9308
		UNIPROT ID:	ENSG00000112149
		Full Name:	CD83 molecule
		Calculated MW:	205 aa, 23 kDa

Applications

Tested Applications:	Positive Controls:
FC	FC : human monocyte-derived mature dendritic cells,
Species Specificity:	human, non-human primates

Background Information

CD83 is a member of the immunoglobulin (Ig) superfamily, consisting of an extracellular V-type Ig-like domain, a transmembrane domain, and a cytoplasmic tail. CD83 is one of the most prominent markers for fully matured dendritic cells (DCs) (PMID: 17334966). It is also found on the surface of other activated hematopoietic cells, including lymphocytes, monocytes, macrophages, and neutrophils (PMID: 31231400; 17334966). CD83 regulates the maturation, activation, and homeostasis of numerous immune cells (PMID: 31231400; 32362900). CD83 can also be expressed as a soluble form. Soluble CD83 can bind to DCs and inhibit their maturation (PMID: 17334966; 12403928).

Storage

Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human monocyte-derived mature dendritic cells were surface stained with 5 μ l Coralite® Plus 488 Anti-Human CD83 (HB15e) Mouse IgG2a RecAb (CL488-65581, Clone: HB15e) (red) or Coralite® Plus 488 Mouse IgG2a Isotype Control (C1.18.4) (CL488-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.