For Research Use Only

CoraLite® Plus 488-conjugated Alpha Tubulin Monoclonal antibody



Catalog Number: CL488-66031

3 Publications

Basic Information

Applications

Catalog Number: CL488-66031 Size:

1000 µg/ml Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG18034

Tested Applications:

human, mouse, rat, Canine

human

GenBank Accession Number:

BC009314 GeneID (NCBI): 10376 **UNIPROT ID:** P68363 Full Name: tubulin, alpha 1b Calculated MW:

50 kDa Observed MW:

50-55 kDa

Purification Method:

Protein A purification

CloneNo.: 1E4C11

Recommended Dilutions:

WB 1:300-1:600 IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

FC (Intra), IF/ICC, WB **Cited Applications:**

Species Specificity:

Cited Species:

Positive Controls:

WB: HepG2 cells, HeLa cells, HEK-293 cells, HSC-T6

cells, NIH/3T3 cells

IF: HeLa cells, mouse embryo tissue

Background Information

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha tubulin is useful for scientists across fields as an internal control due to its high, ubiquitous expression pattern. Tubulin expression $may\ vary\ according\ to\ resistance\ to\ antimic robial\ and\ antimitotic\ drugs.\ This\ anitbody\ is\ CL488 (Ex/Em\ 488\ nm/515)$ nm) conjugated.

Notable Publications

Author	Pubmed ID	Journal	Application
Xueguang Zhang	31571197	Clin Genet	
Chuan Jiang	33968654	Transl Androl Urol	IF
Yaoqiu Wu	36638901	Free Radic Biol Med	IF

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

Selected Validation Data




E14.5 FFPE mouse embryo section stained for smooth muscle actin (red, Cat. No 80008-1-RR) and alpha tubulin (green, Cat. CL488-66031). The alpha tubulin was conjugated to CoraLite-488 fluorescent dye. In red, you can see muscle tissue clearly stained including the tongue, heart, and arteries. Nerves and neurons are stained green since they are rich in tubulin. Image credit:


@Immunofluorescence on Instagram



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using the CoraLite® Plus 488-conjugated version of this antibody, CL488-66031 (alpha Tubulin antibody), at dilution of 1:100.



Various lysates were subjected to SDS PAGE followed by western blot with CL488-66031 (Alpha Tubulin antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human Alpha Tubulin (CL488-66031, Clone:1E4C11) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).