For Research Use Only

CoraLite® Plus 488-conjugated RBAP48 Monoclonal antibody

Catalog Number:CL488-66060

Basic Information	Catalog Number: CL488-66060	GenBank Accession Number: BC053904	Purification Method: Protein A purification
	Size: 1000 µg/ml	GenelD (NCBI): 5928	CloneNo.: 5C4D6
	Source: Mouse	UNIPROT ID: Q09028	Recommended Dilutions: IF/ICC 1:50-1:500
	Isotype: IgG2b Immunogen Catalog Number: AG6196	Full Name: retinoblastoma binding protein 4 Calculated MW: 48 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
		Applications	
Species Specificity: human	IF/ICC : A549 cells,		
Background Information	Histone-binding protein RBBP4 (also known as RbAp48, or NURF 55) is a protein that in humans is encoded by the RBBP4 gene. This gene encodes a ubiquitously expressed nuclear protein that belongs to a highly conserved subfamily of WD-repeat proteins. It is present in protein complexes involved in histone acetylation and chromatin assembly. It is part of the Mi-2/NuRD complex complex that has been implicated in chromatin remodeling and transcriptional repression associated with histone deacetylation. This encoded protein is also part of corepressor complexes, which is an integral component of transcriptional silencing. It is found among several cellular proteins that bind directly to retinoblastoma protein to regulate cell proliferation. A decrease of RbAp48 in the dentate gyrus (DG) of the hippocampus in the brain is suspected to be a main cause of memory loss in normal aging (PMID: 23986399).		
Storage	Storage: Store at -20°C. Avoid exposure to I Storage Buffer:	ight. Stable for one year after shipment	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A549 cells using CoraLite® Plus 488-conjugated RBAP48 antibody (CL488-66060, Clone: 5C4D6) at dilution of 1:200, CL594-Phalloidin (red).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human RBAP48 (CL488-66060, Clone:5C4D6) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL488-66360-3, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).