## For Research Use Only

## CoraLite® Plus 488-conjugated RCAS1 Monoclonal antibody



Catalog Number: CL488-66170

**Basic Information** 

Catalog Number:

CL488-66170

 $\begin{array}{lll} \text{Size:} & \text{GeneID (NCBI):} \\ \text{1000 } \mu \, \text{g/ml} & \text{9166} \\ \text{Source:} & \text{UNIPROT ID:} \end{array}$ 

Mouse 000559
Isotype: Full Name:

G1 estrogen receptor binding site associated, antigen, 9

Immunogen Catalog Number: associated, antiger
AG2905 Calculated MW:

213 aa, 24 kDa Observed MW: 34 kDa

GenBank Accession Number:

Purification Method:

Protein G purification

CloneNo.: 4H8A12

**Recommended Dilutions:** 

IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

Tested Applications: FC (Intra), IF/ICC

Species Specificity:

human

Positive Controls:

IF: MCF-7 cells,

## **Background Information**

Estrogen receptor-binding fragment-associated antigen 9 (EBAG9) gene was identified as an estrogen-responsive gene. The gene product, receptor-binding cancer antigen expressed on SiSo cells (RCAS1), is associated with aggressive characteristics and poor overall survival for 15 different human malignancies. The correlation between RCAS1 expression and several clinicopathological variables, including tumor size, clinical stage, invasion depth and lymph node metastasis highlights this molecule's clinical significance. Expression of RCAS1 in tumor cells plays an important role in evasion from host immune system resulting tumor progression, invasion and metastasis. Further exploration of RCAS1 biological function will facilitate development of novel therapeutic strategies that target RCAS1.

Storage

Storage:


Store at -20 °C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CoraLite® Plus 488 RCAS1 antibody (CL488-66170, Clone: 4H8A12) at dilution of 1:200.



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human RCAS1 (CL488-66170, Clone:4H8A12) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).