For Research Use Only

CoraLite® Plus 488-conjugated P62/SQSTM1 Monoclonal antibody



Catalog Number: CL488-66184

Featured Product

1 Publications

Basic Information

Catalog Number:

CL488-66184 Source:

Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG13131

GenBank Accession Number:

BC017222 GeneID (NCBI):

8878

UNIPROT ID: Q13501

Full Name: sequestosome 1

Calculated MW:

48 kDa
Observed MW:

62 kDa

Purification Method:

Protein A purification

CloneNo.: 1H5C1

Recommended Dilutions:

IF/ICC: 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

IF/ICC

Cited Applications:

IF

Species Specificity:

human
Cited Species:

human

Positive Controls:

IF/ICC: HepG2 cells, Chloroquine treated U2OS cells

Background Information

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and sometimes aggregated proteins for autophagic degradation. As a autophagy-specific substrate, p62 is degraded during the autophagic process, which makes intracellular level of p62 as a marker for autophagy flux. p62 is at the cross-roads of several signaling pathways including Ras/ Raf/ MAPK and NF κ B and plays important role in cancer. p62 is a component of inclusion bodies/ protein aggregates found in human diseases, including Huntington's disease, Alzheimer's disease, Parkinson's disease in the brain, and nephropathic cystinosis in kidney (PMID: 22074114, 22860231, 22714671). The molecular weight of p62 is predicted as 48/ 38 kDa, while western blot analyses using this antibody demonstrate the major band around 60-62 kDa in various tissues.

Notable Publications

Author	Pubmed ID	Journal	Application
Tian-Mei Yu	40460159	PLoS Pathog	IF

Storage

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Chloroquine treated HepG2 cells using CL488-66184 (P62,SQSTM1 antibody) at dilution of 1:100.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL488-66184 (P62,SQSTM1 antibody) at dilution of 1:100.