For Research Use Only

CoraLite® Plus 488-conjugated IDH1 Monoclonal antibody

Catalog Number:CL488-66197




Basic Information	Catalog Number: CL488-66197	GenBank Accession Number: BC012846	Purification Method: Protein G purification
	Size: 1000 ug/ml	GenelD (NCBI): 3417	CloneNo.: 2A6A2
	Source: Mouse	UNIPROT ID: O 75874	Recommended Dilutions: IF/ICC 1:50-1:500
	Isotype: IgG1 Immunogen Catalog Number: AG19293	Full Name: isocitrate dehydrogenase 1 (NADP+), soluble	Excitation/Emission maxima •), wavelengths: 493 nm / 522 nm
		Calculated MW: 414 aa, 47 kDa	
		Observed MW: 46 kDa	
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human	Positive Controls:	
		IF/ICC : HepG	2 cells,
Background Information	IDH1, also named as PICD and IDP, belongs to the isocitrate and isopropylmalate dehydrogenases family. It is a common feature of a major subset of primary human brain cancers. It can form a homodimer(PMID:15173171).IDH1 mutation is always heterozygotic and IDH1 functions as a dimer, theoretically there will be 25% each wild type and mutant homo-dimers and 50% hetero-dimers present in the tumor cells(PMID:21079649).		
Storage	Storage: Store at -20°C. Avoid exposure to l Storage Buffer: PBS with 50% Glycerol, 0.05% Pro Aliquoting is unnecessary for -20°	•	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 IDH1 antibody (CL488-66197, Clone: 2A6A2) at dilution of 1:200.



1x10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488-conjugated IDH1 Monoclonal antibody (CL488-66197, Clone:2A6A2) (red), or 0.4 ug Coralite® Plus 488 Mouse IgG1 Isotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).