For Research Use Only

CoraLite® Plus 488-conjugated INS Monoclonal antibody____

Catalog Number:CL488-66198 2 Publications




Basic Information	Catalog Number: CL488-66198	GenBank Accession Number: BC005255	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	Source:	3630	4B6A7	
	Mouse Isotype:	UNIPROT ID: P01308	Recommended Dilutions: IF-P 1:200-1:800	
	IgG2a	Full Name:	Excitation/Emission maxima	
	- Immunogen Catalog Number:	INS	wavelengths:	
	AG8630	Calculated MW: 110 aa, 12 kDa	493 nm / 522 nm	
Applications	Tested Applications: IF-P	IF-P : mouse pancreas tissue,		
	Cited Applications:			
	Species Specificity:			
	human, mouse, rat			
	Cited Species: mouse			
	INS is a peptide hormone, produced by beta cells of the pancreas, and is central to regulating carbohydrate and fat metabolism in the body. It participates in glucose utilization, protein synthesis and in the formation and storage o neutral lipids. It is synthesized as a precursor molecule, which is processed prior to secretion. A- and B-peptides ar joined together by a disulfide bond to form INS, while the central portion of the precursor molecule is cleaved and released as the C-peptide. Defects in INS results in type 1 diabetes mellitus. INS may also exist 36 kDa form corresponding to the hexameric INS form.			
Background Information	metabolism in the body. It partici neutral lipids. It is synthesized as joined together by a disulfide bor released as the C-peptide. Defects	pates in glucose utilization, protein s a precursor molecule, which is proce Id to form INS, while the central porti s in INS results in type 1 diabetes me	ynthesis and in the formation and storage of ssed prior to secretion. A- and B-peptides and on of the precursor molecule is cleaved and	
	metabolism in the body. It partici neutral lipids. It is synthesized as joined together by a disulfide bor released as the C-peptide. Defect corresponding to the hexameric I	pates in glucose utilization, protein s a precursor molecule, which is proce Id to form INS, while the central porti s in INS results in type 1 diabetes me	ynthesis and in the formation and storage of ssed prior to secretion. A- and B-peptides and on of the precursor molecule is cleaved and	
	metabolism in the body. It partici neutral lipids. It is synthesized as joined together by a disulfide bor released as the C-peptide. Defect corresponding to the hexameric IN	pates in glucose utilization, protein s a precursor molecule, which is proce Id to form INS, while the central porti s in INS results in type 1 diabetes me IS form.	ynthesis and in the formation and storage of ssed prior to secretion. A- and B-peptides ar on of the precursor molecule is cleaved and llitus. INS may also exist 36 kDa form	
Background Information	metabolism in the body. It partici neutral lipids. It is synthesized as joined together by a disulfide bor released as the C-peptide. Defects corresponding to the hexameric IN Author Manxi Yang	pates in glucose utilization, protein s a precursor molecule, which is proce Id to form INS, while the central porti s in INS results in type 1 diabetes me NS form.	ynthesis and in the formation and storage of ssed prior to secretion. A- and B-peptides ar on of the precursor molecule is cleaved and llitus. INS may also exist 36 kDa form Application	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed mouse pancreas tissue using the CoraLite® Plus 488-conjugated version of this antibody, CL488-66198 (INS antibody), at dilution of 1:400. The α cells of pancreas was labelled in red with 15954-1-AP (Glucagon antibody).