## For Research Use Only

## CoraLite® Plus 488-conjugated SETDB1 Monoclonal antibody



Catalog Number: CL488-66293

**Basic Information** 

Catalog Number:

CL488-66293

1000 µg/ml

Source: Mouse Isotype:

lgG1 SET domain, bifurcated 1

Calculated MW: Immunogen Catalog Number:

AG21644 143 kDa

> Observed MW: 170-180 kDa

BC009362

9869

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

Q15047

GenBank Accession Number:

**Purification Method:** 

Protein G purification

CloneNo.: 1H6E5

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human, mouse, rat

## **Background Information**

SETDB1, also named as ESET, KIAA0067 and KMT1E, belongs to the histone-lysine methyltransferase family. It is a SET domain protein with histone H3-K9-specific methyltransferase activity. H3 'Lys-9' trimethylation is coordinated with DNA methylation and represents a specific tag for epigenetic transcriptional repression by recruiting HP1 (CBX1, CBX3 and/or CBX5) proteins to methylated histones. SETDB1 mainly functions in euchromatin regions, thereby playing a central role in the silencing of euchromatic genes. It probably forms a complex with MBD1 and ATF7IP that represses transcription and couples DNA methylation and histone 'Lys-9' trimethylation. Its activity is dependent on MBD1 and is heritably maintained through DNA replication by being recruited by CAF-1. SETDB1 regulates histone methylation, gene silencing, and transcriptional repression. It has been identified as a target for treatment in Huntington Disease, given that gene silencing and transcription dysfunction likely play a role in the disease pathogenesis. The calculated molecular weight of SETDB1 is 143 kDa, but the modified SETDB1 protein is about 170 kDa (PMID: 11791185).


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human SETDB1 (CL488-66293, Clone:1H6E5) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).