For Research Use Only

CoraLite® Plus 488-conjugated SPARC Monoclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Protein A purification

Catalog Number: CL488-66426

Basic Information

Catalog Number: GenBank Accession Number: CL488-66426 BC004974

 Concentration:
 GeneID (NCBI):
 CloneNo.:

 1000 μ g/ml
 6678
 1A2C1

Source: UNIPROT ID: Recommended Dilutions: Mouse P09486 IF/ICC 1:50-1:500

Isotype: Full Name: Excitation/Emission maxima IgG2a secreted protein, acidic, cysteine-rich wavelengths:

Immunogen Catalog Number: (osteonectin) 493 nm / 522 nm

AG7390 Calculated MW:

35 kDa

Observed MW: 38 kDa, 52 kDa

Applications

Tested Applications:

Species Specificity: human, Rat

Positive Controls:

IF/ICC : C6 cells,

Background Information

SPARC, also known as ON (Osteonectin) or BM-40 (Basement-membrane protein 40), is an extracellular glycoprotein with the calculated molecular mass of 35 kDa and the apparent molecular mass of 40-43 kDa and 50 kDa (PMID: 7495300, 12365801). SPARC belongs to a group of matricellular proteins defined as secreted components that do not contribute directly to the formation of structural elements but serve to modulate cell-matrix interactions and cellular functions (PMID: 7542656; 12231357). SPARC is expressed at high levels in bone tissue, is distributed widely in many other tissues and cell types, and is associated generally with tissues undergoing morphogenesis, remodeling and wound repair (PMID: 10567433). It elicits changes in cell shape, inhibits cell-cycle progression, and influences the synthesis of extracellular matrix (PMID: 12721366). Altered expression of SPARC has been reported in a variety of cancers, which include breast, ovarian, colorectal, and pancreatic cancer as well as melanoma and glioblastomas (PMID: 18849185).

Storage

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

torage Buffe


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed C6 cells using Coralite® Plus 488 SPARC antibody (CL488-66426, Clone: 1A2C1) at dilution of 1:200.



1x10^6 A375 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488-conjugated SPARC Monoclonal antibody (CL488-66426, Clone:1A2C1) (red). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).