For Research Use Only

## CoraLite® Plus 488-conjugated STAT5B Monoclonal antibody

Antibodies | ELISA kits | Proteins www.ptglab.com

Catalog Number: CL488-66427

**Basic Information** 

Catalog Number:

CL488-66427

Size: Genel D (NCBI): 1000 ug/ml 6777

 Source:
 ENSEMBL Gene ID:

 Mouse
 ENSG00000173757

 Isotype:
 UNIPROT ID:

 IgG2a
 P51692

Immunogen Catalog Number: Full Name:

AG2709 signal transducer and activator of

Calculated MW: 787 aa, 90 kDa Observed MW: 90 kDa

transcription 5B

GenBank Accession Number:

BC020868

Purification Method:

Protein A purification CloneNo.:

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

1D9B11

**Applications** 

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human

Positive Controls:

IF/ICC: HepG2 cells,

## **Background Information**

STAT5B belongs to the transcription factor STAT family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5B carries out a dual function: signal transduction and activation of transcription. STAT5B mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different GH. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. Despite the 95% homology between the immunogen (Ag2709) and the STAT5A protein, our WB detection showed that this antibody does not recognize the STAT5A fusion protein (Ag19546).

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1x10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488-conjugated STAT5B Monoclonal antibody (CL488-66427, Clone:1D9B11) (red), or 0.4 ug Isotype Control (CL488-66360-1) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 STAT5B antibody (CL488-66427, Clone: 1D9B11) at dilution of 1:200, CL594-Phalloidin (red).