For Research Use Only

CoraLite® Plus 488-conjugated ING2 Monoclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Catalog Number: CL488-66512

Basic Information

Catalog Number: GenBank Accession Number: CL488-66512 BC030128

BC030128 Protein A purification
Genel D (NCBI): CloneNo.:

 $1000 \ \mu \ g/ml$ 3622 2C3A7 Source: UNIPROT ID: Excitation

Source: UNIPROT ID: Excitation/Emission maxima Mouse Q9H160 wavelengths:

 Mouse
 Q9H160
 wavetengths:

 Isotype:
 Full Name:
 493 nm / 522 nm

IgG1 inhibitor of growth family, member 2

Immunogen Catalog Number:Calculated MW:AG19291280 aa, 33 kDa

Observed MW: 33-35 kDa

Applications

Tested Applications:

FC (Intra)

Size:

Species Specificity:

Human, rat, mouse

Background Information

ING2, also named as ING1L, belongs to the ING family and contains a PHD-type zinc finger. As a candidate tumor suppressor, ING2 seems to be involved in p53/TP53 activation and p53/TP53-dependent apoptotic pathways, probably by enhancing acetylation of p53/TP53. It is a component of a mSin3A-like corepressor complex, which is probably involved in deacetylation of nucleosomal histones. Recent finding reported that ING2 regulates muscle differentiation by regulating chormatin remodeling program.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human ING2 (CL488-66512, Clone:2C3A7) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).