For Research Use Only

## CoraLite® Plus 488-conjugated OLIG2 Monoclonal antibody

Isotype:



**Purification Method:** 

Catalog Number: CL488-66513

**Basic Information** 

Catalog Number: GenBank Accession Number: CL488-66513

BC047511 Protein A purification GeneID (NCBI): CloneNo.: 10215 1E8C5

1000 µg/ml **UNIPROT ID:** Recommended Dilutions: Source:

Mouse Q13516 IF 1:50-1:500 Full Name: Excitation/Emission maxima

IgG2a oligodendrocyte lineage transcription wavelengths:

493 nm / 522 nm factor 2 Immunogen Catalog Number:

AG18838 Calculated MW:

32 kDa Observed MW: 35-39 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

Human, mouse, rat

Positive Controls:

IF: mouse brain tissue,

## **Background Information**

OLIG2, also named as BHLHB1, BHLHE19, PRKCBP2 and RACK17, is required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Cooperates with OLIG1, OLIG2 establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced V3 interneuron development. OLIG2 is widely expressed in subsets of glia cells and progenitors, and it is strongly induced at different sites by both acute and chronic injury, albeit with different mechanisms. OLIG2 acts as a repressor of neurogenesis in cells reacting to brain injury. It may represent an effective approach towards evoking neuronal repair from parenchymal precursors. (PMID:19390819 )


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using CoraLite® Plus 488 OLIG2 antibody (CL488-66513, Clone: 1E8C5) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using CoraLite® Plus 488 OLIG2 antibody (CL488-66513, Clone: 1E8C5) at dilution of 1:200.