For Research Use Only

CoraLite®488-conjugated PARP1 Monoclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number:CL488-66520

Basic Information

Catalog Number:
CL488-66520

Size:
1000 μg/ml

Source:
Mouse

Isotype:
IgG1

Immunogen Catalog Number:
AG19173

GenBank Accession Number:
BC037545

GeneID (NCBI):
142

UNIPROT ID:
P09874

Full Name:
poly (ADP-ribose) polymerase 1

Calculated MW:
1014 aa, 113 kDa

Purification Method:
Protein G purification

CloneNo.:
1D7D4

Recommended Dilutions:
IF 1:50-1:500

Excitation/Emission maxima wavelengths:
491 nm / 516 nm

Applications

Tested Applications:
FC (Intra), IF/ICC,IF-P

Species Specificity:
human, mouse

Positive Controls:

IF : human lung cancer tissue, Neuro-2a cells

Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24-kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered as an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and gave rise to fragments ranging from 42-89-kD. This antibody was generated against the N-terminal region of human PARP1 and it recognizes the full-length as well as the cleavage of the PARP1.

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human lung cancer tissue using CoraLite®488 PARP1 antibody (CL488-66520, Clone: 1D7D4) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using CoraLite®488 PARP1 antibody (CL488-66520, Clone: 1D7D4) at dilution of 1:1600.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite®488 Anti-Human PARP1 (CL488-66520, Clone:1D7D4) (red), or 0.2 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).