

For Research Use Only

# CoraLite® Plus 488-conjugated NF- $\kappa$ B p65 Monoclonal antibody



Catalog Number: CL488-66535

Featured Product

## Basic Information

Catalog Number:

CL488-66535

Size:

1000  $\mu$ g/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG1199

GenBank Accession Number:

BC011603

GeneID (NCBI):

5970

UNIPROT ID:

Q04206

Full Name:

v-rel reticuloendotheliosis viral oncogene homolog A (avian)

Calculated MW:

65 kDa

Observed MW:

65 kDa

Purification Method:

Protein G purification

CloneNo.:

1B12D11

Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

## Applications

Tested Applications:

IF/ICC

Species Specificity:

Human, rat

Positive Controls:

IF : HepG2 cells,

## Background Information

Nuclear factor  $\kappa$ B (NF- $\kappa$ B) is a sequence-specific DNA-binding protein complex which regulates the expression of viral genomes, including the human immunodeficiency virus, and a variety of cellular genes, particularly those involved in immune and inflammatory responses. The members of the NF- $\kappa$ B family in mammalian cells include the proto-oncogene c-Rel, p50/p105 (NF $\kappa$ B1), p65 (RelA), p52/p100 (NF $\kappa$ B2), and RelB. All of these proteins share a conserved 300-amino acid region known as the Rel homology domain which is responsible for DNA binding, dimerization, and nuclear translocation of NF- $\kappa$ B. The p65 subunit is a major component of NF- $\kappa$ B complexes and is responsible for trans-activation. NF- $\kappa$ B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF- $\kappa$ B p65-p65 complex appears to be involved in invasion-mediated activation of IL-8 expression. The inhibitory effect of I- $\kappa$ B upon NF- $\kappa$ B in the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF- $\kappa$ B complex. It associates with chromatin at the NF- $\kappa$ B promoter region via association with DDX1.

## Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL488-66535 (NF- $\kappa$ B p65 antibody) at dilution of 1:100.