For Research Use Only

# CoraLite® Plus 488-conjugated SMARCA4 Monoclonal antibody



**Purification Method:** 

Protein A purification

Catalog Number: CL488-66561

**Basic Information** 

Catalog Number: GenBank Accession Number: CL488-66561 BC150298

 Size:
 GeneID (NCBI):
 CloneNo.:

 1000 μg/ml
 6597
 2E6B6

Source: UNIPROT ID: Recommended Dilutions: Mouse P51532 IF/ICC 1:50-1:500

Isotype: Full Name: Excitation/Emission maxima IgG2b SWI/SNF related, matrix associated, wavelengths:

chromatin, subfamily a, member 4

IgG2b SWI/SNF related, matrix associated, wavelengths:
actin dependent regulator of 493 nm / 522 nm

AG16256 chromatin, subfai Calculated MW: 1647 aa, 185 kDa

**Applications** 

Tested Applications: IF/ICC

Species Specificity:

Human

Positive Controls:

IF/ICC: HepG2 cells,

#### **Background Information**

SMARCA4, also named as BAF190A, BRG1, SNF2B and SNF2L4, belongs to the SNF2/RAD54 helicase family. SMARCA4 is a transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. It is a component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. It is also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. And this anitbody is CL488 (Ex/Em 488 nm/515 nm) conjugated.

### Storage

#### Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CL488-66561 (SMARCA4 antibody) at dilution of 1:50.