For Research Use Only

CoraLite® Plus 488-conjugated cIAP1 Monoclonal antibody



Purification Method:

493 nm / 522 nm

Catalog Number: CL488-66626

Basic Information

Catalog Number: GenBank Accession Number: CL488-66626 BC016174

BC016174 Protein G purification
GeneID (NCBI): CloneNo.:

Size: GenelD (NCBI): CloneNo.: 1000 μ g/ml 329 1H3F1

Source: UNIPROT ID: Recommended Dilutions:

Mouse Q13490 IF/ICC 1:50-1:500

Isotype: Full Name: Excitation/Emission maxima

 Isotype:
 Full Name:
 Excitation/En

 IgG1
 baculoviral IAP repeat-containing 2
 wavelengths:

Immunogen Catalog Number: Calculated MW:

AG21398 618 aa, 70 kDa

Applications

Tested Applications:

IF/ICC

Species Specificity: Human, mouse, rat, pig Positive Controls:

IF/ICC : HepG2 cells,

Background Information

BIRC2 (also known as cIAP1) is a member of the inhibitor of apoptosis protein (IAP) family. The inhibitor of apoptosis (IAP) proteins are a family of anti-apoptotic regulators found in viruses and metazoans. BIRC2 is a nuclear shuttling protein, whose subcellular localization is mediated by the CRM1-dependent nuclear export pathway (PMID: 15265700). The protein is regulated transcriptionally and can be inhibited by mitochondrial proteins released in the cytoplasm upon apoptotic stimuli (PMID: 15187025). BIRC2 is also believed to be a critical regulator of vascular integrity and endothelial cell survival, thereby providing an additional target pathway for the control of angiogenesis and blood vessel homeostasis during embryogenesis, regeneration and tumorigenesis (PMID: 17934460).

Storage

Storage:


Store at -20°C. Avoid exposure to light.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 cIAP1 antibody (CL488-66626, Clone: 1H3F1) at dilution of 1:100.