## For Research Use Only

## CoraLite® Plus 488-conjugated NRF1/nuclear respiratory factor 1 Monoclonal antibody




## Catalog Number: CL488-66832

Basic Information	Catalog Number: CL488-66832	GenBank Accession Number: BC016925	Purification Method: Protein A purification				
	Size: 1000 μg/ml	GeneID (NCBI): 4899	CloneNo.: 3C2B8				
	Source: Mouse	UNIPROT ID: Q16656	Recommended Dilutions: IF/ICC 1:50-1:500				
	Isotype: IgG2b Immunogen Catalog Number: AG28149	Full Name: nuclear respiratory factor 1 Calculated MW: 522 aa, 56 kDa Observed MW: 70 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm				
				Applications	Tested Applications:	Positive Controls:	
					IF/ICC, FC (Intra) Species Specificity: human	IF/ICC : H	lepG2 cells,
Background Information	NRF1, also named as Alpha-pal and NRF-1, belongs to the NRF1/Ewg family. It is a transcription factor that activates the expression of the EIF2S1 (EIF2-alpha) gene. NRF1 is implicated in the control of nuclear genes required for respiration, heme biosynthesis, and mitochondrial DNA transcription and replication. It plays a critical role in regulating the expression of many antioxidant response element (ARE)-dependent genes. NRF1 exists various isoforms and long isoform is about 67 kDa and short isoform is about 45 kDa.						
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage						


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 NRF1 antibody (CL488-66832, Clone: 3C2B8) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human NRF1/nuclear respiratory factor 1 (CL488-66832, Clone:3C2B8) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL488-66360-3, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).