For Research Use Only

CoraLite® Plus 488-conjugated NeuN Monoclonal antibody



Catalog Number: CL488-66836 **2 Publications**

Basic Information

Catalog Number: CL488-66836

1000 µg/ml Source: Mouse

lgG1 Immunogen Catalog Number:

AG28016

Isotype:

GenBank Accession Number:

NM_001082575 GeneID (NCBI): 146713

Full Name:

Recommended Dilutions: hexaribonucleotide binding protein 3 IF-P 1:50-1:500

> Excitation/Emission maxima wavelengths:

Purification Method:

Protein G purification

CloneNo.:

3A4C1

493 nm / 522 nm

Applications

Tested Applications: IF-P, FC (Intra) Cited Applications:

IF

Species Specificity: human, mouse, rat **Cited Species:** mouse, rat

Positive Controls:

IF-P: mouse cerebellum tissue, mouse brain tissue

Background Information

NeuN, encoded by FOX3, is a neuron-specific nuclear protein. Anti-NeuN stains exclusively neuronal cells in the central and peripheral nervous systems, especially postmitotic and differentiating neurons, as well as terminally differentiated neurons. Anti-NeuN has been used widely as a reliable tool to detect most postmitotic neuronal cell types. The immunohistochemical staining is primarily localized in the nucleus of the neurons with lighter staining in the cytoplasm.

Notable Publications

Author	Pubmed ID	Journal	Application
Jingcheng Shi	38585980	bioRxiv	
Lirong Zhang	36801237	Exp Eye Res	IF


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using CoraLite® Plus 488 NeuN antibody (CL488-66836, Clone: 3A4C1) at dilution of 1:200.



1X10^6 U-87 MG cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human NeuN (CL488-66836, Clone:3A4C1) (red), or 0.4 ug Mouse IgG1 Isotype Control (66360-1-Ig, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



1X10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human NeuN (CL488-66836, Clone:3A4C1) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).