For Research Use Only

CoraLite® Plus 488-conjugated Hexokinase 2 Monoclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

FC (Intra): 0.40 ug per 10^6 cells in a

WB: 1:1000-1:8000

100 µl suspension

CloneNo.:

2A11C3

Catalog Number: CL488-66974

Basic Information

Catalog Number: GenBank Accession Number: CL488-66974 BC021116

 Source:
 GeneID (NCBI):

 Mouse
 3099

 Isotype:
 UNIPROT ID:

IgG1 P52789
Immunogen Catalog Number: Full Name:

AG16895 hexokinase 2

102 kDa wavelengths: 493 nm / 522 nm Observed MW:

Observed MW: 102 kDa

Calculated MW:

Applications

Tested Applications: WB, FC (Intra) Species Specificity:

human, mouse, rat

Positive Controls:

WB: HeLa cells, HepG2 cells FC (Intra): HepG2 cells,

Background Information

Hexokinase 2 (HK2), a rate-limiting enzyme in the first step of glycolysis pathway, expresses at high level in cancer cells compared with normal cells. HK2 provides a new target for cancer therapy due to its pivotal role in tumor tumourigenic and metastatic process. HK1 is constitutively expressed in most mammalian adult tissues. HK2, however, although is abundantly expressed in embryonic tissues, is expressed at high levels only in limited number of adult tissues such as adipose, skeletal, and cardiac muscles. (PMID: 29305912, PMID: 28427443, PMID: 23911236)

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1x10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Hexokinase 2 Monoclonal Antibody (CL488-66974, Clone:2A11C3)(red), or 0.4 ug Coralite® Plus 488 Mouse IgG1 Isotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.



Various lysates were subjected to SDS PAGE followed by western blot with CL488-66974 (Hexokinase 2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.