

For Research Use Only

CoraLite® Plus 488-conjugated ADC Monoclonal antibody



Catalog Number: **CL488-67884**

Basic Information

| | | |
|---|--|--|
| Catalog Number: CL488-67884 | GenBank Accession Number: BC028128 | Purification Method: Protein G purification |
| Size: 1000 µg/ml | GeneID (NCBI): 113451 | CloneNo.: 1A2B11 |
| Source: Mouse | UNIPROT ID: Q96A70 | Recommended Dilutions: IF/ICC 1:50-1:500 |
| Isotype: IgG1 | Full Name: arginine decarboxylase | Excitation/Emission maxima wavelengths: 493 nm / 522 nm |
| Immunogen Catalog Number: AG31049 | Calculated MW: 460 aa, 50 kDa | |
| | Observed MW: 48-50 kDa | |

Applications

| | |
|---|---|
| Tested Applications: IF/ICC, FC (Intra) | Positive Controls: IF/ICC : A431 cells, |
| Species Specificity: human, mouse, rat, pig | |

Background Information

Antizyme inhibitor 2 (AZIN2) is a member of the antizyme inhibitor family, which is able to bind to AZs and inhibit their action on both ODC activity and polyamine uptake. AZIN2 is predominantly expressed in the brain and testis and localized in the endoplasmic reticulum-golgi intermediate compartment. Available data suggest its role in cell growth, spermiogenesis, vesicular trafficking and secretion. Accumulation of azin2 has also been observed in brains of patients with Alzheimer's disease (PMID: 19832840). AZIN2 has 5 described isoforms produced by alternative splicing. The molecular weight of AZIN2 is about 50 kDa.

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using Coralite® Plus 488 ADC antibody (CL488-67884, Clone: 1A2B11) at dilution of 1:200.



1X10⁶ A431 cells were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human ADC (CL488-67884, Clone:1A2B11) (red), or 0.2 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).