For Research Use Only

CoraLite® Plus 488-conjugated PSME1 Monoclonal antibody



Catalog Number: CL488-67928

Basic Information

Catalog Number: CL488-67928

Size:
1000 µg/ml
Source:
Mouse
Isotype:
IgG1

Immunogen Catalog Number:

AG30549

Observed MW: 29 kDa

Calculated MW: 29 kDa

BC007503

5720

Q06323

GeneID (NCBI):

UNIPROT ID:

Full Name:

GenBank Accession Number:

proteasome (prosome, macropain) activator subunit 1 (PA28 alpha)

Purification Method:

Protein G purification

CloneNo.: 1F7E2

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity: Human, mouse

Background Information

The principal function of the proteasome is targeted degradation of intracellular proteins. Activity of the 20S proteasome is controlled by regulatory complexes that bind to the ends of the cylindrical proteasome. 11S regulator (REG or PA28), is a complex of 28 kDa subunits that is thought to activate proteasomes toward the production of antigenic peptides. Human PSME1 and PSME2 genes encode the two proteasome activators PA28 alpha and beta, respectively, which have been implicated in antigen processing for loading class I MHC molecules. The PA28 activator complex enhances the generation of class I binding peptides by altering the cleavage pattern of the proteasome.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human PSME1 (CL488-67928, Clone:1F7E2) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).