For Research Use Only

CoraLite® Plus 488-conjugated CLN3 Monoclonal antibody

Catalog Number:CL488-67957

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Basic Information

Catalog Number:
CL488-67957

Size:
1000 μg/ml

Source:
Mouse

Isotype:
IgG1

Immunogen Catalog Number:
AG31402

GenBank Accession Number:
BC002394

GeneID (NCBI):
1201

UNIPROT ID:
Q13286

Full Name:
ceroid-lipofuscinosis, neuronal 3

Calculated MW:
438 aa, 48 kDa

Observed MW:
50 kDa

Purification Method:
Protein G purification

CloneNo.:
1E10A9

Excitation/Emission maxima wavelengths:
493 nm / 522 nm

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human

Background Information

Neuronal ceroid lipofuscinosis (NCL, or Batten disease) refers to a group of lethal pediatric neurodegenerative diseases originating from mutations in one of the thus far identified 13 CLN genes (Ceroid Lipofuscinosis, Neuronal type; CLN1 to CLN14) (PMID: 25051496). CLN3 is a multi-membrane spanning protein that is involved in microtubule-dependent, anterograde transport of late endosomes and lysosomes. The CLN3 gene is located on chromosome 16p12.1and produces three mRNA splicing variants. The 438-amino-acid CLN3 protein has a calculated molecular weight of 48 kDa. It has been reported that CLN3 can be glycosylated and form homodimeric complex (PMID: 10356317; 17286803).

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human CLN3 (CL488-67957, Clone:1E10A9) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).