

For Research Use Only

CoraLite® Plus 488-conjugated Phospho-EIF2S1 (Ser51) Monoclonal antibody



Catalog Number: **CL488-68023** **1 Publications**

Basic Information

Catalog Number: CL488-68023	GenBank Accession Number: NM_004094	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 1965	CloneNo.: 1A4A11
Source: Mouse	UNIPROT ID: P05198	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
Isotype: IgG1	Full Name: eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa	
	Calculated MW: 36 kDa	
	Observed MW: 36 kDa	

Applications

Tested Applications:
FC (Intra)

Cited Applications:
IF

Species Specificity:
Human, Rat, Mouse

Background Information

EIF2S1 is one subunit of the translation initiation factor EIF2, which catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B. This phosphorylation stabilizes the eIF2-GDP-eIF2B complex and inhibits the turnover of eIF2B. Induction of PKR by IFN- γ and TNF- α induces potent phosphorylation of eIF2 α at Ser51.

Notable Publications

Author	Pubmed ID	Journal	Application
Yonger Xue	38272900	Nat Commun	IF

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ PC-3 cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug CoraLite® Plus 488 Anti-Human Phospho-EIF2S1 (Ser51) (CL488-68023, Clone:1A4A11), or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.



1X10⁶ PC-3 cells untreated or treated with Calyculin A were intracellularly stained with 0.5 ug CoraLite® Plus 488 Anti-Human Phospho-EIF2S1 (Ser51) (CL488-68023, Clone:1A4A11). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.