For Research Use Only

## CoraLite® Plus 488-conjugated Phospho-STAT5A (Tyr694)/STAT5B (Tyr699) Recombinant antibody



Catalog Number: CL488-80115

**Basic Information** 

Catalog Number: CL488-80115

1000 ug/ml Source: Rabbit

Isotype:

GenBank Accession Number:

BC027036 GeneID (NCBI): 6776 **UNIPROT ID:** P42229 Full Name:

signal transducer and activator of transcription 5A

Calculated MW: 794 aa. 92 kDa Observed MW: 90-95 kDa

**Purification Method:** Protein A purification

CloneNo.: 2P21

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

## **Background Information**

STAT5A, also named as STAT5, belongs to the transcription factor STAT family. In response to cytokines and growth  $factors, STAT\ family\ members\ are\ phosphorylated\ by\ the\ receptor\ associated\ kinases,\ and\ then\ form\ homo-or\ property.$ heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5A is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different GHs. Phosphorylation at Tyr694 is obligatory for Stat5 activation. Activation of STAT5A in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of human STAT5A is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of STAT5A in cells.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 TF-1 cells untreated (dashed lines) or treated with GM-CSF (red) were intracellularly stained with 0.13 ug Coralite® Plus 488 Anti-Human Phospho-STAT5A (Tyr694) (CL488-80115, Clone:2P21), or 0.13 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.