For Research Use Only

CoraLite® Plus 488-conjugated NRF2, NFE2L2 Recombinant antibody

Catalog Number: CL488-80593 Featured Product



Purification Method:

CloneNo.:

1|21

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

wavelengths: 493 nm / 522 nm

Basic Information

Catalog Number: CL488-80593

Size:
1000 ug/ml
Source:
Rabbit
Isotype:

Immunogen Catalog Number:

AG9489

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse

GenBank Accession Number:

BC011558 GeneID (NCBI): 4780 UNIPROT ID:

Q16236

Full Name: nuclear factor (erythroid-derived 2)-

like 2
Calculated MW:

605 aa, 68 kDa Observed MW: 110 kDa

Positive Controls:

IF/ICC: MG132 treated HepG2 cells,

Background Information

Applications

NRF2, also named as NFE2L2, belongs to the bZIP family and CNC subfamily. It is a transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. NRF2 is important for the coordinated up-regulation of genes in response to oxidative stress. It may be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region. Nrf2 is a key player in the regulation of genes encoding for many antioxidative response enzymes. The expression of NRF2 may be induced under oxidative stress (PMID:14567983). In lung cancer, Nrf2 activation in malignant cells has been associated with tumor progression and chemotherapy resistance(PMID:20534738). Identifying patients with abnormal NRF2 expression may be important for selection for chemotherapy in NSCLC. As new investigators break into the emerging field of Nrf2 research, confusion regarding the correct migratory pattern of Nrf2 is causing doubts about the accuracy and reproducibility of published results. This letter provides solid evidence that the actually observed molecular weight of Nrf2 is about 70kDa and 95-110 kDa. (PMID: 22703241).

Storage


Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data







1X10^6 MCF-7 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human NRF2, NFE2L2 (CL488-80593, Clone:1121) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).