For Research Use Only

CoraLite® Plus 488-conjugated MPO Recombinant monoclonal antibody



Catalog Number: CL488-81610 Featured Product

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

CL488-81610

GeneID (NCBI):

CloneNo.:

Source: Rabbit

4353

3J17

Isotype IgG

UNIPROT ID: P05164

Recommended Dilutions:

Immunogen Catalog Number:

Full Name:

IF-P: 1:50-1:500

AG17564

myeloperoxidase

Excitation/Emission maxima wavelengths: 493 nm / 522 nm

Calculated MW: 745 aa, 84 kDa

Observed MW:

59 kDa, 90 kDa

Applications

Tested Applications:

Positive Controls:

Species Specificity:

human

IF-P: human tonsillitis tissue,

Background Information

The MPO gene encodes myeloperoxidase, a lysosomal hemoprotein located in the azurophilic granules of polymorphonuclear (PMN) leukocytes and monocytes. In response to stimulation, MPO is activated into a transient intermediate with potent antimicrobial oxidizing abilities(PMID:17650507). The mRNA is translated into a single protein of 90 kDa, which displays enzymatic activity and undergoes proteolytic maturation into a heavy chain of 59 kDa and a light chain of 13.5 kDa; these subunits then dimerize into the mature tetramer and the mature MPO is a heterotetramer composed of two identical heavy chains and two identical light chains (PMID:12773517). Fragments with molecular masses of 43-47 kDa were formed by autocatalysis during warming in sample buffer (PMID:12960244). The 24-kDa material had a map identical to that of 13.5 kDa subunit and represents a dimer of the 13.5 kDa subunit (PMID:3008892). Defects in MPO are the cause of myeloperoxidase deficiency (MPOD). It has 3 isoforms produced by alternative splicing.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using CoraLite® Plus 488 MPO antibody (CL488-81610, Clone: 3J17) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).