

For Research Use Only

CoraLite® Plus 488-conjugated Caspase 3/P17/19 Recombinant antibody

Catalog Number: CL488-82202



Basic Information

Catalog Number: CL488-82202	GenBank Accession Number: NM_004346	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 836	CloneNo.: 5G20
Source: Rabbit	UNIPROT ID: P42574	Recommended Dilutions: IF/ICC 1:50-1:500
Isotype: IgG	Full Name: caspase 3, apoptosis-related cysteine peptidase	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
	Calculated MW: 32 kDa	
	Observed MW: 32-35 kDa, 17 kDa, 19 kDa	

Applications

Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : HeLa cells,
Species Specificity: human, mouse	

Background Information

Caspases, a family of endoproteases, are critical players in cell regulatory networks controlling inflammation and cell death. Initiator caspases (caspase-2, -8, -9, -10, -11, and -12) cleave and activate downstream effector caspases (caspase-3, -6, and -7), which in turn execute apoptosis by cleaving targeted cellular proteins. Caspase 3 (also named CPP32, SCA-1, and Apopain) proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at the beginning of apoptosis. Caspase 3 plays a key role in the activation of sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase 3 can also form heterocomplex with other proteins and performs the molecular mass of 50-70 kDa. This antibody can recognize p17, p19 and p32 of Caspase 3.


Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 Caspase 3/P17/19 antibody (CL488-82202, Clone: 5G20) at dilution of 1:200.



1X10⁶ HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human Caspase 3/P17/19 (CL488-82202, Clone:5G20) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).