For Research Use Only

CoraLite® Plus 488-conjugated ICAM-2/CD102 Recombinant monoclonal antibody



Catalog Number: CL488-84857-2

Basic Information

Catalog Number: CL488-84857-2

GenBank Accession Number: NM_001099789.2

Purification Method: Protein A purification

Source: Rabbit

GeneID (NCBI): 3384

CloneNo.: 242237F8

Isotype:

UNIPROT ID: P13598

Recommended Dilutions: IF/ICC: 1:50-1:500

Immunogen Catalog Number: EG2002

Full Name: intercellular adhesion molecule 2 Excitation/Emission maxima

Calculated MW:

wavelengths: 493 nm / 522 nm

31kDa

Applications

Tested Applications:

IF/ICC

Positive Controls:

Species Specificity:

human

IF/ICC: Jurkat cells,

Background Information

ICAM2 is a cell adhesion protein having important roles in cell migration, especially during inflammation when leukocytes cross the endothelium. Initially described as a receptor for lymphocyte function-associated antigen-1 (LFA1), ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-recirculation, and other cellular interactions important for immune response and surveillance. ICAM2 has six N-recirculation for immune response and surveillance interactions important for immune response and surveillance interactions in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immune response and surveillance in the six N-recirculation for immunelinked glycosylation sites at amino acids (asparagines) 47, 82, 105, 153, 178 and 187.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using Coralite® Plus 488 ICAM-2/CD102 antibody (CL488-84857-2, Clone: 242237F8) at dilution of 1:200.