For Research Use Only

CoraLite® Plus 488-conjugated SP3 Recombinant antibody

Catalog Number:CL488-85050




Basic Information	Catalog Number: CL488-85050	GenBank Accession Number: BC126414	Purification Method: Protein A purification
	Concentration: 1000 ug/ml	GenelD (NCBI): 6670	CloneNo.: 242556A2
	Source: Rabbit	UNIPROT ID: Q02447	Recommended Dilutions: IF/ICC: 1:200-1:800
	lsotype: IgG	Full Name: Sp3 transcription factor	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
	Immunogen Catalog Number: AG25082	Calculated MW: 781 aa, 82 kDa	
		Observed MW: 70 kDa, 115 kDa	
Applications	Tested Applications:	Positive Controls: IF/ICC : HepG2 cells,	
	Species Specificity:		
	human, mouse, rat		
Background Informatio	 human, mouse, rat SP3, also named as transcription f belongs to the Sp1 C2H2-type zind when sumoylated. Some localizat deacetylated by HDACs and sumo being the major site. Sumoylatior and PML nuclear bodies. Sumoylar isoform, L-Sp3, which has little eff 	c-finger protein family. SP3 localizes t tion in PML nuclear bodies. SP3 is acet bylated on all isoforms. Sumoylated o n at this site promotes nuclear localiza tion on Lys-551 represses the transact fect on transactivation. Alternate sum	, which contains 3 C2H2-type zinc fingers and to the nuclear periphery and in nuclear dots ylated by histone acetyltransferase p300, n 2 sites in longer isoforms with Lys-551 stion to the nuclear periphery, nuclear dots ivation activity, except for the largest oylation and acetylation at Lys-551 also nolecular weight of those are 70 kda and 115
Background Informatio	 human, mouse, rat SP3, also named as transcription f belongs to the Sp1 C2H2-type zind when sumoylated. Some localizat deacetylated by HDACs and sumo being the major site. Sumoylatior and PML nuclear bodies. Sumoylar isoform, L-Sp3, which has little eff control transcriptional activity. Th kda. Storage: 	c-finger protein family. SP3 localizes to tion in PML nuclear bodies. SP3 is acet sylated on all isoforms. Sumoylated on at this site promotes nuclear localize tion on Lys-551 represses the transact fect on transactivation. Alternate sum- here exits tow modified isoform and m light. Stable for one year after shipme	ylated by histone acetyltransferase p300, n 2 sites in longer isoforms with Lys-551 ition to the nuclear periphery, nuclear dots ivation activity, except for the largest oylation and acetylation at Lys-551 also nolecular weight of those are 70 kda and 115

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 SP3 antibody (CL488-85050, Clone: 242556A2) at dilution of 1:400, CL594-Phalloidin (red).