

For Research Use Only

CoraLite® Plus 488 Mouse IgG2a, λ Isotype Control Recombinant Antibody

Catalog Number:CL488-98600



Basic Information

Catalog Number:

CL488-98600

Concentration:

100ug, 500 ug/ml

Source:

Mouse

Isotype:

IgG2a, lambda

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Purification Method:

Protein A purification

CloneNo.:

252253A1

Recommended Dilutions:

FC: 0.25 ug per 10^6 cells in a 100 μ l suspension

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

FC

Species Specificity:

n/a

Positive Controls:

FC : rat splenocytes, human PBMCs

Background Information

This product is an isotype control antibody, which is used to estimate the non-specific binding of target primary antibodies due to Fc receptor binding or other protein-protein interactions. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ rat splenocytes were surface stained with 0.25 ug CoraLite® Plus 488 Mouse IgG2a, λ Isotype Control RecAb (CL488-98600, Clone:252253A1)(red), or 0.25 ug CoraLite® Plus 488 Anti-Rat CD43 (W3/13) Mouse IgG2a Recombinant Antibody (CL488-65607, Clone: W3/13)(green), or unstained (blue). Cells were not fixed.



1x10⁶ human PBMCs were surface stained with 0.25 ug CoraLite® Plus 488 Mouse IgG2a, λ Isotype Control RecAb (CL488-98600, Clone:252253A1)(red), or 0.25 ug CoraLite® Plus 488 Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody (CL488-65559, Clone: UCHT4)(green), or unstained (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.